



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : C12N 15/00, 9/64, A61K 37/54 C12N 1/00	A1	(11) International Publication Number: WO 87/07907 (43) International Publication Date: 30 December 1987 (30.12.87)
(21) International Application Number: PCT/GB87/00420 (22) International Filing Date: 16 June 1987 (16.06.87) (31) Priority Application Numbers: 8614722 8619272 (32) Priority Dates: 17 June 1986 (17.06.86) 7 August 1986 (07.08.86) (33) Priority Country: GB (71) Applicant (for all designated States except US): CELL-TECH LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : DOCHERTY, Andrew, James, Penrose [GB/GB]; 34 Seymour Road, Hampton Hill, Middlesex TW12 1DD (GB). MURPHY, Gillian [GB/GB]; 204 Cambridge Road, Great Shelford, Cambridge CB2 5UV (GB).		(74) Agent: VOTIER, Sidney, David; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS FOR THE PRODUCTION OF A PROTEIN		
(57) Abstract		
<p>Process for the production of a mammalian stromelysin by a recombinant DNA process. The invention also relates to specific proteins, DNA sequences, vectors, host organisms, pharmaceutical compositions, DNA probes, antibodies and to the use of stromelysin to enhance the activity of collagenase. Stromelysin may be useful therapeutically for example in the debridement of dermal ulcers, modification of scar tissue formation arising from healing of wounds and in the treatment of herniated vertebral discs.</p> <div style="text-align: center;"> </div> <p>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 174</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

PROCESS FOR THE PRODUCTION OF A PROTEINField of the Invention

This invention relates to a process for the production of a protein, particularly a metalloproteinase by a recombinant DNA process. The invention also relates to specific proteins, DNA sequences, vectors, host organisms and to pharmaceutical compositions.

Background to the Invention

The metalloproteinases are a family of enzymes produced in mammalian tissues and which are believed to play an important role in the resorption of the extracellular matrices of connective tissues.

The family of metalloproteinases includes collagenase, gelatinase and stromelysin which work together synergistically to digest all the major macromolecules comprising extracellular matrices. The enzymes are found in latent proenzyme form activatable by trypsin or 4-aminophenylmercuric acetate (APMA). The metalloproteinases have a zinc atom at their active site and require calcium for full activity.

Usually tissue metalloproteinases are coordinately synthesised, and for many tissues a specific stimulus is needed to induce synthesis (Reynolds J. Brit. J. Dermatology (1985) 112 715-723). The metalloproteinases are present naturally in low concentration and cannot be extracted in large amounts from mammalian tissues.

Of particular interest is stromelysin which is known to degrade many components of the connective tissues such as proteoglycan core protein, non-helical regions of type IV collagen, laminin, fibronectin, elastin, gelatin and procollagens types I, II and III and which may play a role in biological processes such as wound

healing. Stromelysin may be useful therapeutically in processes such as debridement of dermal ulcers and modifying scar tissue formation arising from the healing of wounds such as burns or necrosis. Stromelysin may also be used in the treatment of herniated vertebral discs to effect dissolution of nucleus pulposus. In order to use stromelysin in this way it must be produced on a large scale and at a commercially viable cost. We provide mammalian stromelysin in such commercially worthwhile amounts by producing it in accordance with the invention, using recombinant DNA techniques.

Preprostromelysin is a large molecular weight form of stromelysin which on secretion is processed to yield the lower molecular weight prostromelysin. Activation of prostromelysin by, for example, treatment with 4-amino phenylmercuric acetate yields the mature biologically active stromelysin.

Summary of the Invention

According to the present invention there is provided a process for the production of a mammalian stromelysin comprising culturing host cells transformed with a DNA sequence coding for the stromelysin.

The term mammalian stromelysin as used herein denotes a mammalian stromelysin having the amino acid sequence of an authentic mammalian stromelysin, an analogue thereof, or a biologically active peptide fragment of either of these, having the biological activity associated with authentic mammalian stromelysin.

In a preferred embodiment the invention provides a process for the production of human stromelysin comprising culturing host cells transformed with a DNA sequence coding for human stromelysin.

The process allows the production of relatively large quantities of stromelysin and, for the first time, facilitates complete characterisation of the structure and pharmacological properties of the protein.

The stromelysin preferably has an amino acid sequence of greater than 90 percent homology (common amino acids/total amino acids) with the sequence of amino acids 1 to 477 of human preprostromelysin as shown in Figure 4 of the accompanying drawings. More preferably, the homology is greater than 98 per cent and most preferably the human preprostromelysin has the amino acid sequence substantially as shown in Figure 4.

In addition to the complete mammalian stromelysin proteins, it is envisaged that relatively small peptide fragments of such proteins e.g. a peptide fragment of stromelysin may be useful as a metalloproteinase. Thus, in a particular embodiment of the first aspect of the invention, and where appropriate in subsequent aspects of the invention, the term stromelysin includes a peptide fragment of a stromelysin protein.

The mammalian stromelysin is suitably a methionine-stromelysin or a methionine-prostromelysin. It is currently understood that to obtain expression of a DNA sequence, the DNA sequence must possess a 5' ATG codon and the corresponding polypeptide therefore possesses an N-terminal methionine amino acid. As used herein the term "methionine-stromelysin" and "methionine-prostromelysin" denote an authentic mammalian stromelysin or prostromelysin, (or an authentic mammalian stromelysin or prostromelysin, modified or substituted to provide a functionally equivalent protein) having an N-terminal methionine residue. Preferably the methionine residue is adjacent to the N-terminal amino acid of the stromelysin or prostromelysin but may be separated therefrom by one or more amino acids provided that the protein possesses stromelysin or prostromelysin functional activity.

In a second aspect of the invention there is provided a process for the production of a mammalian stromelysin comprising producing a precursor of the stromelysin by culturing host cells transformed with a DNA sequence coding for the stromelysin precursor and cleaving the precursor to produce the stromelysin.

In a preferred embodiment of the second aspect of the invention

- 4 -

we provide a process for the production of human stromelysin comprising producing a precursor of human stromelysin by culturing host cells transformed with a DNA sequence coding for the stromelysin precursor and cleaving the precursor to produce human stromelysin.

5 The precursor protein may be a methionine-stromelysin or methionine-prostromelysin.

 The precursor protein may be a mammalian stromelysin or a prostromelysin with an amino terminal signal sequence. The signal sequence may be a sequence having the effect of encouraging
10 transport of the expression products from the host cell in which the DNA sequences have been expressed. For example, in the case of human stromelysin, a signal sequence of amino acids such as depicted by amino acids 1 to 17 in Figure 4 of the accompanying drawings may be attached to the amino terminal amino acid to form
15 preprostromelysin. This signal sequence assists in the export of the product from eukaryotic host cells and is itself cleaved from the product during transport through the cell membrane.

 The precursor protein may be a fusion protein comprising a heterologous protein and a mammalian stromelysin or prostromelysin
20 protein. The heterologous protein may be all or a part of a protein capable of being produced in a host organism, preferably at a high level. Such heterologous proteins include β -galactosidase, chloramphenicol acetyl transferase (CAT) and the product of the trpE gene. The fusion protein preferably includes a site susceptible to
25 selective chemical or enzymatic cleavage between the stromelysin or prostromelysin protein and the heterologous protein. The heterologous protein may be a yeast signal sequence and the host organism may be a yeast. In this preferred embodiment, the yeast host organism advantageously cleaves the fusion protein to produce a
30 mature stromelysin or prostromelysin.

 The process of the invention permits the production of essentially pure human stromelysin, prostromelysin or preprostromelysin. The term essentially pure is used to denote stromelysin which is essentially free from other proteins of human origin.

- 5 -

In a third aspect the invention therefore provides an essentially pure mammalian stromelysin, or stromelysin or preprostromelysin characterised in that the mammalian stromelysin, prostromelysin or preprostromelysin is human stromelysin, prostromelysin or preprostromelysin.

In a preferred embodiment of the third aspect of the invention the essentially pure human stromelysin has substantially the amino acid sequence of from 100 to 477 as shown in Figure 4 of the accompanying drawings, the essentially pure human prostromelysin has the amino acid sequence of from 18 to 477 as shown in Figure 4 of the accompanying drawings and the essentially pure human preprostromelysin has the amino acid sequence of from 1 to 477 as shown in Figure 4 of the accompanying drawings.

In a fourth aspect of the invention, we provide a mammalian stromelysin or prostromelysin produced by the process of the first aspect of the invention or a mammalian stromelysin or prostromelysin precursor protein produced as an intermediate compound.

In a preferred embodiment of the fourth aspect of the invention the stromelysin is human stromelysin or prostromelysin or a precursor thereof.

In a fifth aspect of the invention, we provide a fusion protein comprising a heterologous protein and a mammalian stromelysin or prostromelysin.

In a preferred embodiment of the fifth aspect of the invention the fusion protein comprises a heterologous protein and human stromelysin or human prostromelysin.

In a sixth aspect of the invention, we provide a DNA sequence coding for the amino acid sequence of a mammalian stromelysin or prostromelysin, or a precursor thereof provided that when the DNA sequence codes for rat stromelysin the DNA sequence is not the pTR1 cDNA sequence described by Matrisian et al (EMBO Journal 4 14350-1440 (1985)). Preferably, we provide a DNA sequence coding

for human preprostromelysin having substantially the nucleotide sequence from 51 to 1484 inclusive as shown in Figure 4 of the accompanying drawings. Preferably we provide a DNA sequence coding for human prostromelysin having substantially the nucleotide sequence from 102 to 1484 inclusive as shown in Figure 4 of the accompanying drawings. Preferably we provide a DNA sequence coding for human stromelysin having substantially the nucleotide sequence from 348 to 1484 inclusive as shown in Figure 4 of the accompanying drawings.

10 In a seventh aspect of the invention, we provide an expression vector including a DNA sequence coding for a mammalian stromelysin, prostromelysin or precursor thereof. The vector is adapted for use in a given host cell by the provision of suitable selectable markers, promoters and other control regions as appropriate.

15 In a preferred embodiment of the seventh aspect of the invention the vector includes a DNA sequence coding for human stromelysin or prostromelysin or a precursor thereof.

In an eighth aspect of the invention, we provide host cells transformed with a vector according to the seventh aspect of the invention. The host cells may be any host organism which may be transformed with a vector including a DNA sequence coding for a mammalian stromelysin or a precursor thereof, such that expression of the DNA sequence occurs. Suitable host cells include yeasts (for example Saccharomyces cerevisiae) and mammalian cells in tissue culture (for example, hamster ovary or mouse mammary tumour cells). Preferably, where the host cell is a bacterium or a yeast, the vector includes a DNA sequence coding for methionine-stromelysin or methionine prostromelysin or a fusion protein including a stromelysin or prostromelysin and when the host cell is a mammalian cell in tissue culture, the vector preferably includes a DNA sequence coding for a prestromelysin or a preprostromelysin.

In a ninth aspect of the invention, we provide a pharmaceutical composition comprising a mammalian stromelysin preferably human stromelysin and a pharmaceutically acceptable

- 7 -

excipient. The pharmaceutical composition may be, for example, in the form of an injectable solution or in a form suitable for topical administration. The pharmaceutical composition may contain a stromelysin or a prostromelysin preferably human stromelysin or human prostromelysin which may be activated in vitro or in vivo to give biologically active stromelysin.

In a preferred embodiment of the ninth aspect of the invention we provide a pharmaceutical composition containing an effective amount of a mammalian stromelysin or prostromelysin, preferably human stromelysin or prostromelysin, for use in the debridement of dermal ulcers, modification of scar tissue formation arising from the healing of wounds such as burns and necrosis and in the treatment of herniated vertebral discs.

In a tenth aspect of the invention we provide a process for the production of a pharmaceutical composition comprising bringing a mammalian stromelysin or prostromelysin, preferably human stromelysin or prostromelysin into association with a pharmaceutically acceptable carrier.

In an eleventh aspect the invention provides a method of therapy comprising treating a patient with an effective amount of a mammalian stromelysin preferably human stromelysin, for instance for debridement of dermal ulcers, modification of scar tissue formation arising from the healing of wounds such as burns and necrosis or treatment of herniated vertebral discs.

Preferably, the stromelysin produced in accordance with the invention will have a useful pharmacological effect without significant antigenic reaction with the immune system. In particular the compounds may be used in the debridement of dermal ulcers, modification of scar tissue formation arising from the healing of wounds such as burns and necrosis and in the treatment of herniated vertebral discs.

The DNA sequence coding for human stromelysin as described in Figure 4 may be used to design DNA probes for use in

identifying the over expression of stromelysin indicative of connective tissue disease states and the invention extends to such DNA probes.

5 In a twelvth aspect the invention provides a DNA hybridisation probe comprising a sequence of nucleotides selected from the nucleotide sequence from 1 to 1810 as shown in Figure 4 of the accompanying drawings.

10 In a further aspect the invention provides an antibody having specificity for an antigenic determinant of a mammalian stromelysin. The antibody may be a polyclonal or monoclonal antibody. The antibody may be labelled with a detectable label.

Stromelysin is shown to be able to enhance the activity of collagenase after its activation with, for example, trypsin or APMA and the invention extends to the use of stromelysin for this purpose.

15 Brief Description of the Drawings

The invention is further described by the following non-limiting examples which refer to the accompanying diagrams Figures 1 to 9 and Tables 1 and 2, short particulars of which are given below.

20 Figure 1 shows the N. terminal amino acid sequence of rabbit APMA activated stromelysin (a) and the oligo-nucleotide probe based on this sequence. The 50 base oligonucleotide probe was made in two 25 base halves (b and c) and the complementary 26-mer (d) was synthesised in
25 order to facilitate their ligation. The numbering of the amino acids refers to their position in the cDNA predicted rabbit stromelysin sequence (see Fig 2)

30 Figure 2 Shows the partial nucleotide sequence of rabbit stromelysin and the predicted amino acid sequence

- Figure 3 Shows the restriction maps of 3 human stromelysin cDNAs
- Figure 4 Shows the restriction map and nucleotide sequence of human stromelysin cDNA and the predicted amino acid sequence
- 5
- Figure 5 Shows electrophoretic analysis of the products of activation of stromelysin purified from:
- panel A - human gingival fibroblast culture medium.
- 10 panel B - that secreted by C127 cells transfected with stromelysin-containing vector
- panel C - that secreted by COS cells transfected with stromelysin-containing vector
- Figure 6 Shows a graph of the time course of activation of C127 cell secreted prostromelysin by various concentrations of trypsin or APMA
- 15
- Figure 7 Shows electrophoretic analysis of the products of C127 prostromelysin activation with APMA
- Figure 8 Shows a graph of the effect of different amounts of purified human fibroblast prostromelysin or prostromelysin secreted by C127 cells on the activation by trypsin of:
- 20
- A). procollagenase secreted by transfected C127 cells, and
- 25 B). procollagenase purified from human gingival fibroblast culture
- Figure 9 Shows electrophoretic analysis of procollagenase from human gingival fibroblast culture medium activated by APMA, APMA + stromelysin, trypsin, or trypsin + stromelysin.
- 30

Table 1 Amino acid composition of human prostromelysin

Table 2 Amino acid composition of human stromelysin

Detailed Description of the Embodiments

5 In the following example, prostromelysin is activated by treatment with 4-amino phenylmercuric acetate to yield mature stromelysin. It will be appreciated, however, that by using alternative methods of cleavage of the proenzyme such as enzymic digestion, different forms of active stromelysin may be produced and these are included within the scope of the present invention.

10 1.1 THE ISOLATION OF A RABBIT cDNA ENCODING STROMELYSIN

 4-aminophenylmercuric acetate (APMA) activated (Cawston and
Murphy in: Methods in Enzymology, Volume 80 pp711-722 (1981))
rabbit fibroblast stromelysin was purified from the culture media of
calvariae (Galloway, W. A. et al Biochem. J. 209, 741-752 (1983)).
15 It was reduced and carboxymethylated and subjected to N-terminal
sequencing by automated Edman degradation on an Applied Biosystems
gas-phase sequenator. The 50 base oligonucleotide probe was
designed according to previously described rules to be capable
theoretically of hybridizing to both rabbit stromelysin and
20 collagenase cDNAs (Grantham R. et al Nucleic Acids Res. 9, r43-r47
(1981) and Lathe, R. J. Molec. Biol. 183, 12252-12258 (1983)). It
was synthesized as two continuous 25 mers (b and c, Figure 1) by
automated solid-phase phosphotriester chemistry followed by
purification by HPLC (Patel T. P. et al Nucleic Acids Res. 10,
25 5605-5619 (1982)). A complementary 26 mer (d) was also made (see
Fig. 1).

 Rabbit fibroblasts from explants of normal synovium were grown
to confluence in Dulbecco's modified Eagle's medium (DMEM)
supplemented with 10% fetal calf serum (FCS). After washing they
30 were then maintained in serum-free media for 48 h in the presence of

50-200 ng ml⁻¹ 12 - 0 - tetradecanoylphorbol - 13 - acetate (TPA). mRNA was isolated from approximately 5.10⁷ cells by the guanidinium isothiocyanate/hot phenol method (Maniatis et al 1982 in. Molecular Cloning: A laboratory Manual; Cold Spring Harbor laboratory, New York), followed by oligo (dT)-cellulose column chromatography (Aviv and Leder, 1972 PNAS USA 69 1408-1412). Standard procedures were used to synthesise cDNA (Gubler and Hoffman, 1983 Gene 25 263-269) and a library of 75,000 plaques was established by using EcoRI linkers to ligate the cDNA into λ gt10 essentially as described by Huynh et al in DNA Cloning Vol 1 ed. D. M. Glover, IRL Press p49-78.

The oligonucleotide probe was prepared by first subjecting to a kinase reaction the 25 mer labelled (c) fig 1 and then ligating it to the 25-mer labelled (b) in the presence of the complementary 26-mer (d). The 50-base oligonucleotide probe resulting from the ligation was purified from a denaturing polyacrylamide gel (Maxam A. M & Gilbert W. in Methods in Enzymology, Volume 65, pp 499-560 (1980)) and then kinased to a specific activity of approximately 3.10⁸ Ce cpm μ g⁻¹.

The plaques were transferred to nitrocellulose filters, denatured, and then prehybridised in 5 x SSC, 5 x Denhardt's solution, 50mM NaH₂PO₄, 100 μ gml denatured salmon sperm DNA for 6h at 60°C. Hybridisation was then undertaken for 16h at 40°C in an identical buffer to which had been added 0.1% SDS and the probe DNA at approximately 4ng ml⁻¹. The filters were then washed in 6 X SSC before a final stringent wash (1 X SSC, 0.1% SDS for 15 min at 50°C) followed by autoradiography for 16 h with an intensifying screen. 10 putatively positive plaques were identified of which one gave positive signals after a second screening undertaken as described above. DNA was isolated and the cDNA insert was subcloned into plasmid pSP64 (Melton D. A. et al Nucleic Acids Res. 12, 7035-7056 (1984)). The complete nucleotide sequence of the cDNA was determined by the dideoxy method (Sanger F., Nicklen S. & Coulson A. R. Proc. Natl. Acad. Sci USA 74 5463-5467 (1977)) after subcloning restriction fragments into M13 and transformation into JM101 (Messing J. & Vieira, J. Gene 19 269-276 (1982)).

- 12 -

As shown in fig 2 the cDNA is 532 nucleotides in length and although lacking a poly A tract does have an open reading frame encoding 170 amino acids. Residues 101-127 are identical with the N-terminal amino acid sequence of rabbit stromelysin (Fig 1a) and the amino acid sequence encoded by the cDNA therefore represents the N-terminal third of rabbit stromelysin. This cDNA was therefore used as a probe to identify the human stromelysin cDNA. It was labelled with ^{32}P α dATP and Klenow fragment using the random hexanucleotide method (Feinberg and Vogelstein, 1983, Anal. Biochem, 132 6-13) to a specific activity of 2.10^7 Ce cpm μg^{-1} .

1.2 ISOLATION OF A HUMAN cDNA ENCODING STROMELYSIN

Human fibroblasts derived from gingival explants (passage 3 to 6) were cultivated in DMEM supplemented with 0.2% lactalbumin hydrolysate and 5% partially purified pig IL-1. After 48h mRNA was isolated from approximately 5×10^8 cells by the guanidinium isothiocyanate/CsCl method (Maniatis *et al* 1982 in Molecular Cloning: A laboratory Manual; Cold Spring Harbor Laboratory, New York). The mRNA was used to generate a library of 200,000 recombinant plaques in $\lambda\text{gt}10$ essentially as described above. The plaques were transferred to nitrocellulose, denatured, and then prehybridised in 6 X SSC, 50% formamide, 5 X Denhardt's solution, $100 \mu\text{g ml}^{-1}$ denatured salmon sperm DNA for 16 h at 42°C . They were then hybridised for 48h at 42°C in an identical buffer to which had been added 0.1% SDS and the rabbit stromelysin probe cDNA at a final concentration of approximately 17 ng ml^{-1} . The filters were then washed with the highest stringency being 2 X SSC, 0.1% SDS at 42°C for 30 minutes before autoradiography with an intensifying screen for 48h. 21 putatively positive plaques were identified of which 16 gave a clear positive signal after a second screening undertaken as described above. The cDNA insert of representative plaques was isolated and subcloned into plasmid pSP64 (Melton D. A. *et al* Nucleic Acids Res. 12, 7035-7056 (1984)). Restriction mapping of the largest inserts confirmed that they were related overlapping DNA segments (Fig. 3) and their nucleotide sequence was compiled by the dideoxy method (Sanger F., Nicklen S. &

Coulson A. R. Proc. Natl. Acad. Sci USA 74 5463-5467 (1977)) after subcloning restriction fragments into M13 and transformation into JM101 (Messing J. & Vieira, J. Gen. 19, 269-276 (1982)). This information is shown in Fig 4, where the solid arrows indicate direction of dideoxy sequencing and the broken arrows show directions of sequencing by the Maxam and Gilbert method (Maxam, A.M. and Gilbert, W., Methods Enzymol. (1980) 65 499-560).

1.3 IDENTIFICATION OF RAT cDNA ENCODING STROMELYSIN

We have discovered a high degree of homology between rabbit stromelysin and another amino acid sequence predicted from a rat cDNA, referred to as pTR1 (Matrisian L. M. et al EMBO Journal 4, 1435-1440 (1985)). This cDNA corresponds to a mRNA which is highly enriched in normal fibroblasts after infection with either polyoma virus or rous sarcoma virus or transfection with either the middle t oncogene or the cellular oncogene H-ras. The same mRNA was also reported to be specifically induced after exposure of fibroblasts to EGF (Matrisian L. M. et al EMBO Journal 4, 1435-1440 (1985)). The predicted Mr of the rat protein encoded by pTR1 is approximately 53000, in close agreement with the in vitro translation product of rabbit stromelysin mRNA (Frisch S. M., Chin J. R. & Werb Z. J. Cell Biol. 97, (2, Pt. 5): 430a (Abstr.) (1983)), and data not shown]. We conclude therefore, that the protein encoded by pTR1 is rat stromelysin. The identification of pTR1 as rat stromelysin allows, for the first time, the production of rat stromelysin using recombinant DNA techniques.

2. NUCLEOTIDE SEQUENCE OF HUMAN STROMELYSIN cDNA AND THE PREDICTED AMINO ACID SEQUENCE

The cDNA is 1825 nucleotides in length and appears to be complete at the 3' end because it has a poly A tail preceded by the polyadenylation signal AATAAA (Proudfoot and Brownlee, 1981, Nature, 252 359-362). It contains an open reading frame (nucleotides 51 to 1481) which translates into a polypeptide of 477 amino acids. Evidence that the cDNA encodes stromelysin comes from the finding that 24 of the 27 amino acids encoded by nucleotides 348 to 428 are

identical with and in the same position as the 27 amino acids identified by N-terminal sequencing the APMA activated rabbit stromelysin (Fig 1a and Fig 4). Furthermore most of the amino acids in the N-terminal third of the molecule are identical with and in the same position as those amino acids predicted from the rabbit partial stromelysin cDNA.

Hydrophobicity plots suggest that human stromelysin is rather water soluble except for a hydrophobic N-terminal sequence of 17 amino acids. This is consistent with these amino acids being the hydrophobic core of a signal sequence which is cleaved during secretion to liberate mature protein. Since it is thought that the metalloproteinases are secreted as proenzymes (Harris E. D. *et al* Collagen Rel Res 4 493-512 (1984)) we conclude that the amino acids found between the signal sequence and the phenylalanine (F) at the N-termini of the APMA - activated form of rabbit stromelysin (residue 100 in the human stromelysin sequence) are cleaved off during activation. Preprostromelysin is therefore considered to be comprised of amino acids 1-477, prostromelysin is considered to be comprised of amino acids 18-477 and stromelysin is considered to be comprised of amino acids 100-477. The amino acid composition of prostromelysin and stromelysin together with the molecular weight of the core proteins is shown in tables 1 and 2 respectively. Potential glycosylation sites within the sequences exist, and it is envisaged that expression of these proteins in appropriate cells (see below) may give rise to glycosylated products.

3. THE PRODUCTION OF STROMELYSIN

3.1 EXPRESSION IN E. COLI

The expression of stromelysin or prostromelysin with an additional NH₂-terminal methionine residue (met-stromelysin or met-prostromelysin) in *E. coli* can be achieved by using oligonucleotide linkers to join the mature stromelysin or prostromelysin encoding sequence described in Section 2 above and shown in Figure 4 with a promoter, Shine Dalgarno sequence and an initiating ATG codon. The *E. coli* *trpE* promoter and Shine Dalgarno sequence may be used.

TABLE 1Amino Acid Composition of Human Prostromelysin

	<u>Residue</u>	<u>Number</u>	<u>%</u>	<u>Weight</u>	<u>%</u>
	F	32	6.96	4704	9.02
	L	38	8.26	4294	8.23
5	I	20	4.35	2260	4.33
	M	6	1.30	786	1.51
	V	28	6.09	2772	5.31
	S	27	5.87	2349	4.50
	P	36	7.83	3492	6.69
10	T	27	5.87	2727	5.23
	A	27	5.87	1917	3.68
	Y	17	3.70	2771	5.31
	H	13	2.83	1781	3.41
	Q	9	1.96	1152	2.21
15	N	16	3.48	1824	3.50
	K	32	6.96	4096	7.85
	D	36	7.83	4140	7.94
	E	30	6.52	3870	7.42
	C	3	0.65	309	0.59
20	W	8	1.74	1488	2.85
	R	23	5.00	3588	6.88
	G	32	6.96	1824	3.50
	TOTAL	460		52162	

TABLE 2Amino Acid Composition of Human Stromelysin

	<u>Residue</u>	<u>Number</u>	<u>%</u>	<u>Weight</u>	<u>%</u>
5	F	30	7.94	4410	10.31
	L	30	7.94	3390	7.92
	I	19	5.03	2147	5.02
	M	3	0.79	393	0.92
	V	19	5.03	1881	4.40
10	S	24	6.35	2088	4.88
	P	32	8.47	3104	7.25
	T	24	6.35	2424	5.67
	A	25	6.61	1775	4.15
	Y	13	3.44	2119	4.95
15	H	12	3.17	1644	3.84
	Q	6	1.59	768	1.79
	N	14	3.70	1596	3.73
	K	22	5.82	2816	6.58
	D	28	7.41	3220	7.53
20	E	25	6.61	3225	7.54
	C	2	0.53	206	0.48
	W	8	2.12	1488	3.48
	R	17	4.50	2652	6.20
	G	25	6.61	1425	3.33
TOTAL		378		42789	

These sequences are present for example on plasmid pCT54 into which DNA sequences encoding proteins of interest can be inserted in a Clal site (Emtage et al (1983), Proc. Nat. Acad. Sci. USA, 80, 3671-3675). However, other plasmids also harbouring these or other promoter and Shine Dalgarno sequences but which are amplifiable and from which expression can be tightly regulated may also be used (see, for example, European patent application EP-A2-0121386). The stromelysin or prostromelysin coding sequence without the signal coding sequence but with an additional N-terminal methionine codon, may be inserted into the E. coli expression vector pMG196. This may be achieved by standard recombinant DNA techniques (Maniatis et al (1982), In: Molecular Cloning: A Laboratory Manual, 390-433). Expression of stromelysin or prostromelysin may also be achieved using analogous techniques to those described for expression of TIMP in European patent application No. 86300042.8.

3.2 PURIFICATION OF STROMELYSIN FROM E. COLI CELLS

E. coli cells harbouring a plasmid constructed as described above, when grown under optimal conditions for expression, will produce met-stromelysin or met-prostromelysin at levels of up to 10% of total cellular protein.

The soluble protein fraction from crude E. coli extracts is assayed for stromelysin activity essentially as described by Galaway et al (1983) Biochem. J. 209 741-752. A part or whole of the met-stromelysin expressed in E. coli may be in an insoluble form and therefore not detected in the above assay. In such circumstances, it is solubilised and activated prior to assay and purification. One example of how this can be achieved in relation to methionine-prochymosin production is described in our co-pending International patent application PCT/GB 83/00152 (published as WO 83/04418) and in published British patent application GB 2100737A. Having obtained soluble, active, partially purified stromelysin it is further purified using standard protein purification techniques, as described for example by Galaway et al (1983) Biochem. J. 209 741-752.

3.3 EXPRESSION IN YEAST

Using standard recombinant DNA techniques in a similar manner to those used to achieve expression in E. coli, plasmid vectors suitable for expression of stromelysin in yeast are constructed.

5 The constructions are based, for example, on the vectors described in co-pending published European patent application EP-A2-0073635. They contain the stromelysin or prostromelysin encoding sequence flanked by the yeast phosphoglycerate kinase (PGK) promoter and the PGK gene 3' untranslated end. The orientation of the stromelysin or

10 prostromelysin cDNA with respect to the PGK promoter is such that it ensures expression of mature stromelysin or prostromelysin with an additional NH₂-terminal methionine residue as described for expression in E. coli. Alternatively, pre-stromelysin or prostromelysin may be expressed by joining the stromelysin or

15 prostromelysin encoding sequence to the vector such that the stromelysin signal sequence is left in place. Expression of a fusion protein between a yeast signal sequence (for example, the yeast α -Factor signal sequence (Kurjan and Herskowitz, (1982), Cell, 30, 933-943), and mature stromelysin or prostromelysin can be

20 achieved through the use of appropriate linkers. These plasmid DNAs are introduced into yeast cells, for example, by the method of Beggs (Nature, (1978), 275, 104-109).

3.4 PURIFICATION OF STROMELYSIN FROM YEAST CELLS

Yeast cells containing these plasmids, when grown under optimal

25 conditions for stromelysin expression, will produce up to 5% of total cellular protein as stromelysin or prostromelysin. Depending on the alternatives described above, stromelysin or prostromelysin may or may not be secreted from the yeast cells. Expressed stromelysin is quantified, assayed and purified essentially as

30 described above if produced intercellularly. If secreted it is purified from cell supernatants by standard protein purification techniques (Galaway et al (1983) Biochem. J. 209 741-752).

3.5 EXPRESSION OF STROMELYSIN IN CULTURED ANIMAL CELLS

The preprostromelysin encoding cDNA was inserted into the different kinds of expression vector designed specifically for: 1. transient expression in COS cells (Gluzman, Y. (1981) Cell 23 175-182) and 2. Stable expression in C127 cells (Lowy, D.R. et al J. Virol. (1978) 26 291-298). The necessary DNA manipulations employed standard techniques (Maniatis et al In. Molecular Cloning, A Laboratory Manual, 1982, Cold Spring Harbor Laboratory, New York). The cDNA was tailored for insertion into the vectors by first converting the Ava I site encoded by nucleotides 1591-1596 (Figure 4) into an EcoR1 site. This was achieved by cleaving the cDNA with Ava I, filling in with T4 polymerase followed by addition of an EcoR1 linker. On cleavage with EcoR1 a 1600 bp EcoR1 fragment encoding preprostromelysin extending from the 5' EcoR1 site encoded by nucleotides 1-6 in Figure 4 to the newly created EcoR1 site is generated. This fragment was purified from a 1% low gelling temperature agarose gel and then ligated between the SV40 late promoter and the SV40 early polyadenylation regulatory elements in a "poison minus" derivative of a pBR322 COS cell vector (Lusky, M and Botchan, M.R., Nature, 1981, 293 79-81). Vector DNA containing the preprostromelysin sequence in the appropriate orientation with respect to the SV40 late promoter was purified from cesium chloride gradients and transfected into COS cells using DEAE dextran (Lopata, M.A. et al Nucleic Acids Res., 1984, 12 5707-5717). Serum-free supernatants were harvested 72 hours after transfection.

For expression in C127 cells the preprostromelysin sequence with the early SV40 polyadenylation regulatory elements was removed on a PstI to BamHI fragment from the COS cell vector and put under the control of the mouse metallothionein I promoter in a bovine papilloma virus (BPV) based vector. This arrangement was essentially as described for the expression of tissue inhibitor of metalloproteinase (TIMP), Docherty et al Nature 1985, 318 66-69 and UK Patent Application GB 2169295A. The BPV vector encoding preprostromelysin was introduced into C127 cells by the calcium phosphate coprecipitation method (Wigler, M. et al 1978, Cell 14 725-731) and CdCl₂ (20μM) - and ZnCl₂ (20μM) - resistant

foci were selected. Stromelysin producing cell lines were identified after 21 days and used to generate serum-free culture medium.

4. STROMELYSIN ACTIVATION, ASSAY AND ELECTROPHORETIC ANALYSIS

5 The stromelysin produced as described above was activated with either trypsin or APMA. Trypsin was incubated with the stromelysin at ratios such as 1:100 to 1:1 respectively followed by the addition of a ten-fold excess of soya-bean trypsin inhibitor. Concentrations of trypsin such as 0.1 to 10 $\mu\text{g ml}^{-1}$ were employed and the
10 temperatures and time of incubation were as described below. Treatment with 4-aminophenylmercuric acetate (APMA) was at 1 to 2 mM for various times as indicated below, at 37°C. Stromelysin activity was assayed using ^{14}C -acetylated casein (Galloway, W.A., et al Biochem. J. 1983, 209 741-752) at 35°C for 1 or 4 h. 1 unit
15 of stromelysin degrades 1 μg of casein min^{-1} at 37°C.

 Stromelysin before and after activation as described above and in more detail below was treated with 20mM EDTA, reduced with 500mM 2-mercaptoethanol and run on 10% polyacrylamide mini gels in the presence of SDS. Gels were electro-transferred to nitrocellulose
20 and the enzyme bands visualised using an antisera against rabbit stromelysin which was able to detect human stromelysin (Murphy, G. et al Collagen Rel. Res. 1986 6 351-364) and a peroxidase-labelled second antibody (Hembry, R.M., et al J. Cell Sci. 1985 73 105-119).

 As shown in Figure 5, prostromelysin preparations A, purified
25 from human gingival fibroblast culture medium, B, secreted by C127 cells transfected with the stromelysin-containing vector and C, secreted by COS cells transfected with the stromelysin-containing vector, were treated as follows: 1. no treatment; 2. incubation with 1mM APMA for 2h at 37°C; 3. incubation with 10 $\mu\text{g/ml}$ trypsin
30 for 30 min at 37°C; 4. as 3. with addition of soya bean trypsin inhibitor after 30 min and further incubation for 2h at 37°C; 5. incubation with 100 $\mu\text{g/ml}$ trypsin for 10 min at 4°C; 6. incubation with 10 $\mu\text{g/ml}$ trypsin for 5 min at 4°C;

7. as 6. with addition of soya bean trypsin inhibitor after 5 min and further incubation for 2h at 37°C. The samples were electrophoresed on a 10% polyacrylamide gel in SDS and reducing conditions, electro-blotted onto nitrocellulose and visualised using a sheep anti rabbit stromelysin antibody, and a peroxidase labelled rabbit anti sheep IgG.

As shown in Figure 6 prostromelysin secreted by C127 cells transfected with the stromelysin-containing vector was incubated with A, (●) 0.1µg/ml trypsin, (Δ) 1µg/ml trypsin, B, (□) 10µg/ml trypsin, C, (V) 100µg/ml trypsin at 37°C (closed symbols) or 4°C (open symbols) in a volume of 20µl, or D, (◆) with 2mM APMA at 37°C. Activity elicited was assayed by degradation of ¹⁴C-casein in a 1h assay at 37°C, after addition of soya bean trypsin inhibitor. Stromelysin incubated without trypsin or APMA had no activity.

Prostromelysin was activated with APMA (as described in Figure 6) for varying lengths of time. The samples were then treated with 20mM EDTA and electrophoresed under reducing conditions on a 10% polyacrylamide gel containing SDS and electroblotted and visualised as described in Fig. 5 and Methods. Figure 7 shows the SDS analysis where Track 1. 0 min; Track 2. 30 min; Track 3. 1h; Track 4. 2h; Track 5. 4h; Track 6. 15h.

The pro-forms of either purified natural stromelysin from human gingival fibroblasts, or the recombinant enzyme secreted by COS or C127 cells had an identical Mr of 57000, with a minor 60000 component (probably due to glycosylation (Nagase, H. et al (1983) Biochem. J. 214 281-288); Figure 5, lanes A1, B1, C1). These forms were completely inactive against casein or other substrates (Galloway, W.A. et al (1983) Biochem. J. 209 741-752) but could be activated by trypsin, (e.g. C127 stromelysin, Fig. 6A-C) or less efficiently by APMA, to degrade these substrates (Figure 6D). Trypsin activation was optimal over a wide range of concentrations; activation could be effected at 4°C, with marginally slower kinetics (Figure 6A-C). Only very high levels of trypsin inactivated stromelysin. Activation by APMA was slower, achieving 58% (COS) to 90% (C127 and

natural) of the trypsin value. Although maximum APMA activation was attained in 4-8 h at 37°C (Figure 6), it was found that activation was dependent upon stromelysin concentration (73% of optimal at a ten fold dilution and 33% at a one hundred fold dilution of a 50 unit/ml preparation). Hence, very low concentrations of enzyme, such as the COS cell culture media were probably not maximally activated even after 15 h incubation.

Analysis of the effect of these treatments on the prostromelysin by gel electrophoresis and immuno blotting showed that a reduction in Mr occurred to yield two major species of Mr. 50000 and 48000 slowly in the case of APMA (e.g. C127 stromelysin, Figure 7) and very rapidly in the case of trypsin (Figure 5 A3; B3,5; C3,5). At longer incubation times with APMA traces of an Mr. 28000 form of stromelysin were generated (Figure 7). Plasmin could also be used to generate bands of the same Mr as those produced by trypsin (data not shown). It was noted that optimal activation (as documented in Figure 6) did not require complete conversion of the upper doublet to the lower doublet (Figure 5; lanes A6, A7; Figure 3, lane 4). Using collagenase, Stricklin *et al* (Biochemistry 22 63-68 (1983)) have reported similar observations with activity detectable prior to an Mr loss. It has been proposed that conformational changes in the intact molecule occur, leading to subsequent self-cleavage. The activity of stromelysin elicited was 360 units/mg.

5. COLLAGENASE ACTIVATION ELECTROPHORETIC ANALYSIS AND ASSAY

The collagenase was activated as described for stromelysin or as detailed below. Electrophoretic analysis was undertaken as described for stromelysin except that enzyme bands were visualised using anti-human collagenase antisera followed by a peroxidase-labelled second antibody (Hembry, R.M. *et al* J. Cell Sci. 1985, 73 105-119). The collagenase was assayed by the ¹⁴C-acetylated collagen diffuse fibril assay at 35°C for 4 hours (Whitham, S.E. *et al* Biochem. J. 1986, 240 913-916). 1 unit of collagenase degrades 1µg of type I collagen min⁻¹ at 35°C.

Figure 8 shows the results when procollagenase preparations, A, secreted by C127 cells transfected with the collagenase containing vector and B, purified from human gingival fibroblast culture medium were incubated with 10µg/ml trypsin at 37°C for 30 min in the presence of varying amounts of (●) purified human fibroblast prostromelysin or (■) prostromelysin secreted by C127 cells transfected with the stromelysin-containing vector, followed by the addition of excess soya bean trypsin inhibitor. Open symbols denote the effect of the corresponding activated stromelysin in the absence of trypsin. Activity was assayed using ¹⁴C-type I collagen at 35°C for 4h. Results are expressed as the fold increase elicited by stromelysin relative to the activity with trypsin alone.

Figure 9 shows electrophoretic analysis of procollagenase purified from human gingival fibroblast culture medium treated as follows: 1. no treatment; 2. incubation with 1mM APMA at 37°C for 2h; 3. as 2, in the presence of purified stromelysin; 4. incubation with 10µg/ml⁻¹ trypsin at 37°C for 30 min, followed by addition of soya bean trypsin inhibitor; 5. as 4. in the presence of stromelysin.

Purified natural procollagenase has an Mr of 55000 with a minor component of Mr 59000 upon analysis by SDS polyacrylamide gel electrophoresis and blotting with a specific antiserum to collagenase (Figure 9; lane 1). The procollagenase could be activated by either APMA or trypsin treatment behaving precisely as described by Stricklin *et al* (Biochemistry 22 63-68 (1983)) with a fall in Mr of 10,000 (Figure 9, lanes 2 and 4). The activity elicited was very low, in the region of 320 units/mg. A similar observation was made by Vater *et al* (J. Biol. Chem. 258 9374-9382 (1983)) for rabbit procollagenase. However, the inclusion of either purified or recombinant human prostromelysin in the activation mixture enhanced the collagenase activity by up to 8 fold (Figure 8B). Similar effects of stromelysin on the molecular weight and activity of recombinant C127 cell derived human collagenase were also obtained. With the recombinant collagenase the stromelysin brought about a 12 fold enhancement in activity (Figure 8A).

5 Addition of previously activated stromelysin to the collagenase after trypsin treatment had less effect on the final collagenase activity detectable (data not shown). Active stromelysin alone elicited similar collagenase activities to those with trypsin alone. The ratio of stromelysin:collagenase for efficient activation was in the order of an excess of 2moles or more per mole. It is possible that higher specific activity stromelysin would be a more efficient activator.


10 Analysis of the changes occurring during these treatments showed that the procollagenase, underwent stepwise reductions in Mr APMA and trypsin generated two bands of about 50000 and 45000 (Figure 9, lanes 2 and 4) which were converted to bands of 48000 and 43000 in the presence of stromelysin (Figure 9; lanes 3 and 5). The same band patterns were generated by both natural and
15 recombinant stromelysin. Activated stromelysin alone appeared to produce very small changes in procollagenase of Mr about 500, with limited further conversion to 48000 and 43000.

CLAIMS

1. A process for the production of a mammalian stromelysin comprising culturing host cells transformed with a DNA sequence coding for the stromelysin.
- 5 2. A process for the production of a mammalian stromelysin comprising producing a precursor of the stromelysin by culturing host cells transformed with a DNA sequence coding for the stromelysin precursor and cleaving the precursor to produce the stromelysin.
- 10 3. An essentially pure mammalian stromelysin, prostromelysin or preprostromelysin characterised in that the mammalian stromelysin, prostromelysin or preprostromelysin is human stromelysin, prostromelysin or preprostromelysin.
- 4 A mammalian stromelysin or prostromelysin produced by a process according to Claim 1.
- 15 5. A fusion protein comprising a heterologous protein and a mammalian stromelysin or prostromelysin.
6. A DNA sequence coding for the amino acid sequence of a mammalian stromelysin or prostromelysin or a precursor thereof provided that when the DNA sequence codes for rat stromelysin the
20 DNA sequence is not the pTR1 cDNA sequence.
7. An expression vector including a DNA sequence coding for a mammalian stromelysin, prostromelysin or precursor thereof.
8. Host cells transformed with a vector according to Claim 7.
9. A pharmaceutical composition comprising a mammalian stromelysin
25 and a pharmaceutically acceptable excipient.
10. A method of therapy comprising treating a patient with an effective amount of a mammalian stromelysin.

2/7

Fig. 3



() = other sites for these enzymes
are on map but not shown

X = XbaI
M = MboII
B = BglII
H = HindIII
N = NcoI
P = PvuII
A = AclI
S = SalI

1. The invention relates to a method for the identification of a specific DNA sequence in a library of DNA fragments.

3/7




Fig. 4

SUBSTITUTE SHEET

4/7




Fig. 5





Fig. 9

SUBSTITUTE SHEET

5/7

Fig. 6



6/7




Fig. 7

SUBSTITUTE SHEET






Fig. 8

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 87/00420

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁴ : C 12 N 15/00; C 12 N 9/64; A 61 K 37/54; C 12 N-1/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁴	C 12 N A 61 K	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	The Journal of Biological Chemistry, volume 260, no. 22, 5 October 1985, The American Society of Biological Chemists, Inc., (US), J.R. Chin et al.: "Stromelysin, a connective tissue-degrading metallo- endopeptidase secreted by stimulated rabbit synovial fibroblasts in parallel with collagenase", pages 12367-12376 see the whole document	3,4 1,2,6-8
Y	--	
X	Biochem. J., volume 209, 1983, The Bio- chemical Society, (GB), W.A. Galloway et al.: "Purification and characterization of a rabbit bone metalloproteinase that degrades proteoglycan and other connective- tissue components", pages 741-752 see the whole document	3,4 1,2,6-8
Y	cited in the application	
	--	
	./.	
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"Z" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
6th October 1987		19 NOV 1987
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		M. VAN MOL 

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	The EMBO Journal, volume 4, no. 6, 1985, IRL Press Limited, (Oxford, GB), L.M. Matrisian et al.: "Epidermal growth factor and oncogenes induce transcription of the same cellular mRNA in rat fibroblasts", pages 1435- 1440 see the whole document cited in the application	1,2,6-8
X	J. Cell Biol., volume 97, no. 2, part 5, 1983, S.M. Frisch et al.: "Molecular cloning of a cDNA encoding a secreted connective tissue-degrading metallo- proteinase induced by changes in cytoskeletal structure", page 460a, abstract 1740 see the whole abstract cited in the application	6 1,2,7,8
P,X	Biochem. J., volume 240, 1986, (GB), S.E. Whitham et al.: "Comparison of human stromelysin and collagenase by cloning and sequence analysis", pages 913-916 see the whole document	6
Y	EP, A, 0157604 (SANKYO CO., LTD) 9 October 1985 see the whole document	1,2,6-8
P,X	Journal of Cell Biology, volume 103, no. 5, part 2, 1986, (US), E.J. Clark et al.: "Synthesis and secretion of collagenase and stromelysin in SV40 transformed human ataxia-telangiectasia fibroblasts", page 257a, abstract 961 see the whole abstract	6
A	Chemical Abstracts, volume 105, no. 17, 27 October 1986, (Columbus, Ohio, US), S.M. Frisch: "Molecular biology of the stromelysin and collagenase genes", see page 316, abstract 148779t, & Diss. Abstr. Int. B 1986, 46(5), 1544	
A	Chemical Abstracts, volume 105, no. 3, 21 July 1986, (Columbus, Ohio, US), G.I. Goldberg et al.: "Human fibro- blast collagenase. Complete primary structure and homology to an oncogene transformation-induced rat protein", ./. .	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

	see page 209, abstract 19690h, & J. Biol. Chem. 1986, 261(14), 6600-5	
P,A	GB, A, 2182665 (MONSANTO CO.) 20 May 1987	9,10
A	EP, A, 0115974 (INSTITUT PASTEUR) 15 August 1984 see the whole document	
A	EP, A, 0159497 (K.K. YAKULT HONSHA) 30 October 1985 see the whole document	9,10

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 10... because they relate to subject matter not required to be searched by this Authority, namely:

See PCT rule 39.1 (IV) Methods for treatment of the human or animal
body by surgery or therapy, as well as
diagnostic methods.

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON

INTERNATIONAL APPLICATION NO. PCT/GB 87/00420 (SA 17576)

This Annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 22/10/87

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0157604	09/10/85	JP-A- 60207583	19/10/85
GB-A- 2182665	20/05/87	None	
EP-A- 0115974	15/08/84	WO-A- 8402653	19/07/84
		FR-A- 2538702	06/07/84
		AU-A- 2410584	02/08/84
		JP-T- 60500413	28/03/85
		FR-A, B 2538701	06/07/84
EP-A- 0159497	30/10/85	JP-A- 60188066	25/09/85
		US-A- 4624924	25/11/86

For more details about this annex :
see Official Journal of the European Patent Office, No. 12/82